

**Effect of Pentachlorophenol on Testicular Steroidogenesis
during Different Reproductive Phases of the Catfish,
*Heteropneustes fossilis***

*Surya Prakash Verma**

*Sunanda Dubey***

Abstract

Endocrine disruptor chemical may result in decline of fish population and affect other animals and human beings after entering into food chain. This study is important in order to generate baseline data for its toxicity in fish and to make an inference needed to evaluate the toxic potency of Pentachlorophenol (PCP). During pre-spawning phase, testes of PCP exposed group (32 µg/l/day for 28 days) showed a decreased in the values of testosterone while an increase in the E2 level was observed which indicates estrogenic potential of PCP whereas, in spawning phase, testes of PCP exposed group showed a significant increase in the testosterone level and decrease in the E2 level. Increase in Estradiol level was found in testes of exposed fish during pre-spawning phase might be due to the effect of PCP as endocrine disruptor and estrogenic as well. Increase in Testosterone level during spawning phase could be due to increased spermatogenesis during this phase. Study with dose and duration dependent exposure may explain further understanding.

Introduction

Pentachlorophenol (PCP) was extensively used for decades in agriculture and industry (Zheng et al., 2012). It is an organic chlorinated compound used worldwide for preserving utility poles, railroad, and as insecticides, pesticides, and biocides for controlling agricultural and household pests (Cooper and Jones, 2008). Deleterious effects of PCP such as developmental toxicity, liver defects, genetic toxicity, and endocrine disrupting activity have been studied in livestock (Morales et al., 2014). The pollution of PCP may pose a severe threat to the survival of aquatic species, and it is crucial to analyze its ecological risk. The adverse effects of PCP have been reported by many workers which included immunotoxicity, carcinogenicity, oxidative stress and metabolic disorders (Chen et al., 2015). The PCP has been demonstrated to display estrogenic, anti-estrogenic and anti-androgenic in vitro

* Assistant Professor, Department of Science, Acharya Sitaram Chaturvedi Mahila Mahavidyalaya, Varanasi-221012, India.

** Department of Zoology, Udai Pratap College, Varanasi-221002, India.

Orton et al., 2009; and disturbance in the hypothalamic-pituitary-gonadal (HPG) axis in vivo study Zhang et al., 2008). The proteins in HPG axis often showed inconsistent values in response to the change of time and concentration of exogenous toxicant Liu et al., 2009). There are studies lacking on the distribution of steroid hormones which play important role in regulating reproductive status and spawning behavior (Bruton, 1996). The PCP has been reported to cause DNA damage, endocrine disruption (Zhang et al., 2014), impairments of ovaries, follicular atresia (Sawle et al., 2010), morphological deformities (Cheng et al., 2015), altered activities of antioxidant enzymes, changes in serum testosterone (Zhang et al., 2014), anti-estrogenicity (Zhao et al., 2006), reproductive organ damage (Zhang et al., 2014), immunotoxicity (Shelley et al., 2009) and mutation (Yin et al., 2009) in fish species. Many endocrine disruptors inhibit and/or induce sex steroid biosynthesis consequently linked to reproductive abnormalities in animals Reeder et al., 2005). Testicular steroids can induce spermiation and certain steroid glucuronides function as sex pheromones implying importance in breeding and captive reproduction (Chaube, et al 2018). Fish is used widely as a bio-indicator of water pollution due to rapid responses with high sensitivity to changes in their usual physiological functions such as molecular, biochemical, cellular, hormonal, or behavioral responses (van der Oost et al., 2003). The freshwater catfish, *H. fossilis* is an edible, economically important fish,

and is ideal for wastewater aquaculture. The present study will be useful in understanding the impact of PCP on metabolic and reproductive functions encountered by freshwater fishes. Steroids are chief hormonal messengers synthesized mainly by gonads and play key role in sexual differentiation, metabolism, osmoregulation and germ cell growth, maturation and release, these are degraded largely in the liver to form water soluble metabolites, which are excreted out (Chaube, et al., 2018). The objective of this study was to investigate possible impact of PCP on steroid profile of testes of the catfish, *H. fossilis*. The present analysis can provide valuable insights with regard to the functions of the steroids in the male fish exposed to the PCP.

Materials and Methods

Chemicals

PCP (Crystalline, 99% pure) was purchased from Acros Organics (Geel, Belgium). All other chemicals were of analytical grade and purchased locally. The PCP was dissolved in ethanol and then diluted with water to obtain the required concentrations 32 µg/l (1/10th of LC₅₀).

Animal collection and maintenance

The experiments were performed in accordance with local/national guidelines for experimentation in animals and all care was taken to prevent cruelty of any kind. The freshwater catfish *H. fossilis* were purchased from the local market in Varanasi and used for the experiment. They were acclimated to laboratory conditions for one week prior to the experiments. They were maintained in the laboratory under normal photo period (13.0L:11.0D) and temperature ($25\pm 2^{\circ}\text{C}$) until use for experiments. The fish were fed with boiled egg daily *ad libitum*.

Experimental setup

The acclimatized adult male fish classified into two groups (15 fish per each): first group control, second group PCP treated (28 days with $32\ \mu\text{g/l/day}$) in pre-spawning and spawning phase. In the present study, the amount of PCP exposures was $32\ \mu\text{g/l}$ and the exposure concentrations were determined by performing LC_{50} experiment (Dubey and Verma, 2019, 2021; Dubey et. al., 2022). The conditions of the experiment were as that of acclimatization with changing all the containers water and concentrations of PCP every day.

Steroid extraction

The tissues were homogenized separately in 4 volume cold PBS (0.02 M, phosphate buffered-saline pH 7.4) with an ultrasonic homogenizer at 0°C for 5–10 seconds. The homogenate was centrifuged at 5000 rpm for 20 min at 4°C and extracted with 3 volume diethyl ether, three times. The ether phase was pooled, evaporated and dried under N_2 gas, and stored at -20°C . The incubation medium was directly extracted with diethyl-ether, as described above. The ether phase was pooled group-wise, evaporated, dried under N_2 and stored at -20°C . For the hormone assay, both tissue supernatant and corresponding incubation medium were pooled to make a sample. The supernatant was collected and steroid extracted in 3 volume diethyl ether, 3 times. Organic fractions were pooled, evaporated and dried under N_2 gas and stored at -20°C till further

estimation. Prepared the extracted steroid in solution form by adding $100\ \mu\text{l}$ methanol to steroid extract.

A. Testosterone assay

Testosterone was assayed using an ELISA kit according to the manufacturer's instructions. Briefly, $25\ \mu\text{l}$ each of standard (0, 20, 100, 400 and $1600\ \text{pg/ml}$) and samples were pipetted into the anti-TIgG-coated plate well. The immunoreaction was started by adding $100\ \mu\text{l}$ of T-HRP conjugate solution to each well, followed by incubation at 37°C for 1 hour. The content from each plate was removed and washed with $300\ \mu\text{l}$ of distilled water, 5–6 times. Water was completely drained out from each

well. Next 100 μ l of 3, 3', 5, 5'-tetramethylbenzidine (TMB) substrate was dispensed into each well and incubated at 25°C for 15 min in dark. Color development was stopped by adding 100 μ l of stop solution (0.15 M, sulphuric acid). Absorbance was taken at 450 nm using a Multiscan microplate reader (Thermo Electron Corporation, USA).

B. Estradiol-17 β (E2) assay

E2 was assayed using an ELISA kit according to the manufacturer's instructions. Briefly, 25 μ l each of standard (0, 20, 120, 300, 600 and 2000 pg/ml) and samples were dispensed into the anti-E2-IgG-coated plate well. The immunoreaction was started by adding 200 μ l of E2-HRP conjugate solution to each well, followed by incubation at 37°C for 2 hour. The content from each plate was removed and washed with 300 μ l of distilled water, 5–6 times. Water was completely drained out from each well. Next 100 μ l of 3, 3', 5, 5'-tetramethyl benzidine substrate was dispensed into each well and incubated at 25°C for 30 min in dark. Color development was stopped by adding 100 μ l of stop solution (0.15 M, sulphuric acid). Absorbance was taken at 450 nm using a Multiscan microplate reader (Thermo Electron Corporation, USA).

Statistical Analysis

The data were expressed as means \pm standard error mean (SEM). Comparisons of means (control and treated fish) were done by Student's t-test. The result was considered significant at 5% level ($p < 0.05$).




Fig. 1: Estimation of Testosterone (T) and estradiol-17 β (E2) in the testes of the catfish *H. fossilis* exposed to 32 μ g/l/ day PCP for 28 days during the pre-spawning phase of reproduction. (Asterisk shows significant difference from Control at $P < 0.05$; Student's t-Test) Data were expressed as mean \pm SEM




Fig. 2: Estimation of Testosterone (T) and estradiol-17 β (E2) in the testes of the catfish *H. fossilis* exposed to 32 μ g/l/ day PCP for 28 days during the spawning phase of reproduction. (Asterisk shows significant difference from Control at $P < 0.05$; Student's t-Test). Data were expressed as mean \pm SEM.

Results

During pre-spawning phase, testes of PCP exposed group (28 days with 32 μ g/l/day) showed a decrease in the values of testosterone while an increase in the E2 level was observed. In spawning phase, testes of PCP exposed group (28 days with 32 μ g/l/day) showed a significant increase in the testosterone level while decrease in the E2 level was observed.

Discussion

Amongst vertebrates, fishes are the most at the risk of endocrine disruption since their habitat receives the greatest input of pollutants due to anthropogenic activities or natural weathering processes. Fishes are, therefore, considered the most suitable animal model in endocrine disruption-related research. To date most of the studies elucidating toxicity of xenobiotics on fishes focus on two of the major endocrine axes, the stress and reproductive (Hontela, 1998). It is reported that sex steroids are synthesized in the brain, renal and gonadal tissue of teleosts (Petersen et al., 2015). Further in studies it is revealed that androgen biosynthesis is under the control of the HPG axis which perceives endogenous and environmental cues to modulate sex steroid synthesis from cholesterol (Miller and Bose, 2011). Steroidogenesis produces the estrogen, E2, and the androgen (Carnevali et al., 2018). The steroidogenic pathway is highly conserved across vertebrate taxa and is a major target of endocrine disruption (Feswick et al., 2014). Initial spermatogenesis is under FSH control, while later spermiogenic stages are regulated by pituitary LH (Itoh et al., 1988). Some studies relate reproductive cycle in adult fish with fluctuating levels of androgen synthesis

(Weltzien et al., 2002). Seasonality was reported in plasma testosterone levels (Borg, 1994) and is directly correlated to spermatogenesis (Rinchard et al., 2001). During late spermatogenesis, a reduction in testosterone is attributed to a shift in the steroidogenic pathway towards 17,20 β -P or similar Mullerian inhibiting hormone (MIH) production (Yaron and Levavi-Sivan, 2011). Androgens also have roles in lipid transport and oxidation (Hoffman et al., 2008), β lymphocyte differentiation (Moens, et al., 2007), xenobiotic clearance (Martyniuk and Denslow, 2012) and protein metabolism (Dorts et al., 2009). Androgens also regulate endothelial function, inflammation and oxidative stress in fishes (Si et al., 2014), although the specific mechanisms through which they accomplish this, is currently unknown.

Several xenobiotic compounds that are reported to disrupt various physiological activities fall in the list of endocrine-disrupting chemicals since minute alteration in hormone levels, their receptors or post-receptor signals can cause significant changes in the functions of target cells and tissues (Colborn et al., 1993).

The PCP is highly persistent with the potential to modulate several biological processes that have an impact on growth and development. It is often detected in the aquatic environment (Crosby et al., 1981). Wu et al (2001) showed that in channel catfish testes, estrogens, participate in the regulation of male gamete development and fertility. In vertebrates ranging from sharks to mammals, testicular tissues have been shown to contain aromatase, the rate-limiting enzyme for estrogen formation (Gist et al., 2007).

In the present study testosterone levels was high in control group in pre spawning phase which was expected. Similar findings state that androgen levels are high moderately during the initiation of spermatogenesis (Rolland et al., 2009). Androgens of the testes increased during spermatogenesis and spermiation and play significant role in male reproduction (Amer et al., 2001).

Two androgen receptor (AR) subtypes (α and β) have been demonstrated in fish and they are all predominantly expressed in the gonad particularly expressed in sertoli and interstitial cells (Ikeuchi et al., 2001) suggesting that androgens develop biological activity via the testicular somatic cells. Similarly, the investigation also showed high levels of 17, 20 β -DP in testes, levels being higher in pre spawning phase in testes than spawning phase. Baynes and Scott (1985) proposed that a major function of 17, 20 β -DP in the male is to control sperm motility mediated by changes in K⁺ composition of seminal fluid. Another function of 17, 20 β -DP is that it can be conjugated into glucuronide and conjugates and these water soluble conjugates have pheromonal role (Ikeuchi et al., 2001). In the catfish 17, 20 β -DP may be involved in sperm maturation and spermiation and role in pheromone is a field to be investigated further. Steroid levels are generally low during the non-reproductive period, but increase gradually throughout gametogenesis and decline abruptly thereafter. The predominance of T, 11-

ketotestosterone (11-KT) and E2 in initiating and regulating seasonal reproductive events are most intensively studied (Zhou et al., 2007).

In catfish *H. fossilis* high E2 in testes in spawning phase was observed. Regarding the functional role of E2, it may promote or inhibit early spermatogonia renewal.

Song and Gutzeit, 2003). The E2 may regulate the regulation of the steroidogenesis (i.e. Star, 3 β -HSD, aromatase A and B). Administration of E2 to maturing sea bream males resulted in the identification of numerous estrogen-dependent genes in the testes (Patino et al., 2006).

In spawning phase, testes of PCP exposed fish showed a significant increase in the testosterone level while a decrease in the E2 level. Increase in testosterone level during spawning phase might be due to increased spermatogenesis during this phase. Study with dose and duration dependent exposure is needed for its further understanding.

Endocrine disruption by PCP was monitored in adult zebrafish and rare minnow (*Gobiocypris rarus*) with elevated plasma thyroxine concentrations (Zha et al., 2007). Many synthetic chemicals have been shown to function as natural steroid hormones since they can interact with the estrogen receptor (ER) as agonists and elicit biological responses (Jin et al., 2012). For example, these endocrine disrupters can bind to ER α or ER β and subsequently alter the normal expression of estrogen-responsive genes (Verderame et al., 2011). In zebrafish and rare minnow, the estrogen receptor (ER) mRNA was up-regulated in males and down-regulated in females, and vitellogenin mRNA induction and serum vitellogenin protein increase was reported in longer exposed females (Zhang et al., 2014). During pre-spawning phase, testes of PCP exposed *H. fossilis* showed a decrease in the values of testosterone and increase in estradiol level, could be due to the effect of PCP as endocrine disruptor and its estrogenic potential as well.

Conclusion

At present time, surface waters are polluting by various ways. Elevated levels of pollutants in aquatic systems have resulted from a number of activities including agriculture, urbanization, impoundments, mining and industrial activities. These effects reduced growth rates, impaired reproduction and sometimes death. Bioaccumulation and bio concentration of these in the food chain can put consumers, including humans at risk. PCP is an organic chlorinated compound used worldwide for preserving utility poles, railroad, and as insecticides, pesticides, and biocides for controlling agricultural and household pests (Cooper and Jones, 2008). During pre-spawning phase, testes of PCP exposed fish showed a decreased in the values of testosterone while an increase in the E2 level was observed which indicates estrogenic potential of PCP. In spawning phase, testes of PCP exposed group showed a significant increase in the testosterone level while decrease in the E2 level. The pollution of PCP

may pose a severe threat to the survival of aquatic species due to its endocrine disruptive activity.

References

- Amer, M.A., Miura, T., Miura, C., & Yamauchi, K. (2001). Involvement of sex steroid hormones in the early stages of spermatogenesis in Japanese huchen (*Hucho perryi*). *Biol Repro*, 65(4): 1057- 1066. <https://doi.org/10.1095/biolreprod65.4.1057> Corpus ID: 188504
- Baynes, S.M., and Scott, A.P. (1985). Seasonal variations in parameters of milt production and in plasma concentrations of sex steroids of male rainbow trout (*Salmo gairdneri*). *Gen Comp Endocrinol*, 57(1): 150-160.
- Borg, B. (1994). Androgens in teleost fishes. *Comp Biochem Physiol C Toxicol Pharmacol*, 109: 219-45.
- Bruton, M.N. (1996). Alternative Life–history strategies of catfishes: In the Biology and culture of catfishes. *Aquat Living Resour*, 9: 35-41.
- Carnevali, O., Santangeli, S., Forner-Piquer, I., Basili, D., & Maradonna, F. (2018). Endocrinedisrupting chemicals in aquatic environment: what are the risks for fish gametes? *Fish Physiol Biochem*, 44(6): 1561-1576.
- Chaube, R. Singh, S., Yadav, S., & D. Kumar, D. (2025). Toxicity of Pentachlorophenol Exposure on Male and Female Heteropneustes fossilis Investigated Using NMR-Based Metabolomics Approach. *ACS Omega*, 10 (7): 6279-7477.
- Chaube, R., Mishra, S., & Singh, R. K. (2018). A comparison of steroid profiles in the testis and seminal vesicle of the catfish (*Heteropneustes fossilis*). *Theriogenology*, 105: 90–96.
- Chen, H.M., Lee, Y.H., & Wang, Y.J. (2015). Ros-triggered signaling pathways involved in the cytotoxicity and tumor promotion effects of pentachlorophenol and tetrachlorohydroquinone. *Chem Res Toxicol*, 28(3): 339-350.
- Cheng, Y., Ekker, M., & Chan, H. (2015). Relative developmental toxicities of pentachloroanisole and pentachlorophenol in a zebrafish model (*Danio rerio*). *Ecotoxicology & Environmental Safety*, 112: 7–14.
- Colborn, T., vom Saal, F.S. & Soto, A.M. (1993). Developmental effects of endocrine disrupting chemicals in wildlife and humans. *Environ Health Perspect*, 101(5): 378–384.
- Cooper, G.S., & Jones, S. (2008). Pentachlorophenol and cancer risk: focusing the lens on specific Chlorophenols and contaminants. *Environmental Health Perspectives*, 116(8): 1001-1008.
- Crosby, D.G., Beynon, K.I., Greve, P.A., Korte, F., Still, G.G., & Vouk, J.W. (1981). Environmental chemistry of penta-chlorophenol. *Pure appl Chem*, 53: 1051-1080.
- Dorts, J., Bauwin, A., Kestemont, P., Jolly, S., Sanchez, W., & Silvestre, F. (2012).

- Proteasome and antioxidant responses in *Cottus gobio* during a combined exposure to heat stress and cadmium. *Comp Biochem Physiol C Toxicol Pharmacol*, 155(2): 318–324.
- Dubey, S., & Verma, S.P. (2019). Histopathological changes in liver of *Heteropneustes fossilis* exposed to Pentachlorophenol (PCP). *Proceedings of Zoological Society of India*, 18(2): 103-108.
 - Dubey, S., & Verma, S.P. (2021). Impact of Pentachlorophenol on histopathological changes in the testes of Catfish, *Heteropneustes fossilis*. *Proceedings of Zoological Society of India*, 20(1): 25-30.
 - Dubey, S., Verma, S.P. & Dwivedi, S. (2022). Estimation of antioxidative enzymes in the testes and liver of the *Heteropneustes fossilis* exposed to pentachlorophenol. *Plant Archives*, 22(1): 95 - 101 .
<http://dx.doi.org/10.51470/PLANTARCHIVES.2022.v22.no1.014>.
 - Feswick, A., Ings, J.S., Doyle, M.A., Bosker, T., Munkittrick, K.R., & Martyniuk, C.J. (2014). Transcriptomics profiling and steroid production in mummichog (*Fundulus heteroclitus*) testes after treatment with 5 α -dihydrotestosterone. *Gen Comp Endocrinol*, 203: 106–119.
 - Gist, D.H., Bradshaw, S., Morrow, C.M.K., Congdon, J.D., & Hess, R.A. (2007). Estrogen response system in the reproductive tract of the male turtle: An immunocytochemical study. *Gen Comp Endocrinol*, 151(1): 27–33.
 - Hontela, A. (1998). Interrenal dysfunction in fish from contaminated sites: *in vivo* and *in vitro* assessment. *Environ Toxicol Chem*, 17(1): 44–48.
 - Ikeuchi, T., Todo, T., Kobayashi, T., Nagahama, Y. (2001). Two subtypes of androgen and progesterone receptors in fish testes. *Comp Biochem Physiol*, 129(2-3): 449–455.
 - Itoh, H., Suzuki, K., & Kawachi, H. (1988). The complete amino acid sequences of beta-subunits of two distinct chum salmon GTHs. *Gen Comp Endocrinol*, 71(3): 438-51.
 - Jin, X., Zha, J., Xu, Y., Giesy, J., & Wang, Z. (2012). Toxicity of pentachlorophenol to native aquatic species in the Yangtze River. *Environmental Science and Pollution Research*, 19(3): 609-618.
 - Luo, J., Lei, B., Ma, M., Zha, J., & Wang, Z. (2011). Identification of estrogen receptor agonists in sediments from Wenyu River, Beijing, China. *Water Research*, 45(13): 3908–3914.
 - Martyniuk, C.J., & Denslow, N.D. (2012). Exploring androgen-regulated pathways in teleost fish using transcriptomics and proteomics. *Integr Comp Biol*, 52(5): 695–704.
 - Miller, W.L., & Bose H.S. (2011). Early steps in steroidogenesis: intracellular cholesterol trafficking. *J Lipid Res*, 52(12): 2111-2135.
 - Moens, L.N., van der Ven, K., Van Remortel, P., Del-Favero, J., & De Coen, W.M. (2007). Gene expression analysis of estrogenic compounds in the liver of common

- carp (*Cyprinus carpio*) using a custom cDNA microarray. *J Biochem Mol Toxicol*, 21(5): 299–311.
- Morales, M., Martinez-Paz, P., Martín, R., Planello, R., Urien, J., Martinez-Guitarte, J.L., & Morcillo, G. (2014). Transcriptional changes induced by in vivo exposure to pentachlorophenol (PCP) in *Chironomus riparius* (Diptera) aquatic larvae. *Aquatic Toxicology*, 157: 1–9.
 - Orton, F., Lutz, I., Kloas, W., & Routledge, E.J. (2009). Endocrine disrupting effects of herbicides and pentachlorophenol: in vitro and in vivo evidence. *Environ Sci Technol*, 43(6): 2144-2150.
 - Patino, P.I.S., Teodosio, H.R., Galay-Borgos, M., Power, D.M., & Sweeney, G.E. (2006). Identification of estrogen-responsive genes in the testis of sea bream (*Sparus auratus*) using suppression subtractive hybridization. *Mol Reprod Dev*, 73(3): 318-329.
 - Petersen, L.H., Hala, D., Cantu, C.M., Martinovic, D., & Huggett, D.B. (2015). Effects of progesterone and norethindrone on female fathead minnow (*Pimephales promelas*) steroidogenesis. *Environ Toxicol Chem*, 34(2): 379-390.
 - Reeder, A.L., Ruiz, M.O., Pessier, A., Brown, L.E., Levengood, J.M., Phillips, C.A., Wheeler, M.B., Warner, R.E., & Beasley, V.R. (2005). Intersexuality and the cricket frog decline: historic and geographic trends. *Environ Health Perspect*, 113(3): 261–265.
 - Rinchard, J., Dabrowski, K., & Ottorbe, J. (2001). Sex steroids in plasma of lake whitefish *Coregonus clupeaformis* during spawning in Lake Erie. *Comp Biochem Physiol C Toxicol Pharmacol*, 129(1): 65-74.
 - Rolland, A.D., Lareyre, J.J., Goupil, A.S., Montfort, J., Ricordel, M.J., Esquerré, D., Hugot, K., Houlgate, R., Chalmel, F., & Le Gac, F. (2009). Expression profiling of rainbow trout testis development identifies evolutionary conserved genes involved in spermatogenesis. *BMC Genomics*, 10:546.
 - Sawle, A.D., Wit, E., Whale, G., & Cossins, A.R. (2010). An information-rich alternative, chemicals testing strategy using a high definition toxicogenomics and zebrafish (*Danio rerio*) embryos. *Toxicological Sciences*, 118(1): 128–139.
 - Shelley, L., Balfry, S., Ross, P., & Kennedy, C. (2009). Immuno toxicological effects of a sub-chronic exposure to selected current-use pesticides in rainbow trout (*Oncorhynchus mykiss*). *Aquatic Toxicology*, 92(2): 95-103.
 - Si, D., Li, J., Liu, J., Wang, X., Wei, Z., Tian, Q., Wang, H., & Liu, G. (2014). Progesterone protects blood-brain barrier function and improves neurological outcome following traumatic brain injury in rats. *Exp Ther Med*, 8(3): 1010–1014.
 - Song, M., & Gutzeit, H.O. (2003). Effect of 17 α -ethynylestradiol on germ cell proliferation in organ and primary culture of medaka (*Oryzias latipes*) testis. *Dev Growth Different*, 45(4): 327-337.

- Van der Oost, R., Beyer, J., & Vermeulen, N.P. (2003). Fish bioaccumulation and biomarkers in environmental risk assessment: a review. *Environmental Toxicology & Pharmacology*, 13(2): 57-149.
- Weltzien, F.A., Taranger, G.L., Karlsen, O., & Norberg, B. (2002). Spermatogenesis and related plasma androgen levels in Atlantic halibut (*Hippoglossus hippoglossus* L.). *Comp Biochem Physiol A Mol Integr Physiol*, 132(3): 567-75.
- Wu, C., Patino, R., Davis, K.B., & Chang, X. (2001). Localization of estrogen receptor alpha and beta RNA in germinal and nongerminal epithelia of the channel catfish testis. *Gen Comp Endocrinol*, 124(1):12-20.
- Yaron, Z., & Levavi-Sivan, B. (2011). Endocrine regulation of fish reproduction. In: A.P. Farrell (Ed), *Encyclopedia of fish physiology: vol. 2. From genome to environment* (pp. 1500-1508). San Diego: Academic Press.
- Yin, D., Zhu, H., Hu, P., & Zhao, Q. (2009). Genotoxic effect of 2, 4, 6-trichlorophenol on P53 gene in zebrafish liver. *Environmental Toxicology & Chemistry*, 28(3): 603-608.
- Zha, J., Wang, Z., Wang, N., & Ingersoll, C. (2007). Histological alternation and vitellogenin induction in adult rare minnow (*Gobiocypris rarus*) after exposure tethynylestradiol and nonylphenol. *Chemosphere*, 66(3): 488-495.
- Zhang, M., Yin, D., & Kong, F. (2008). The changes of serum testosterone level and hepatic microsomal enzyme activity of crucian carp (*Carassius carassius*) exposed to a sublethal dosage of pentachlorophenol. *Ecotox Environ Safe*, 71(2): 384-389.
- Zhang, X., Zhang, X., Qi, Y., Huang, D., & Zhang, Y. (2014). 2, 4-dichlorophenol induces ER stress-mediated apoptosis via eIF2 α dephosphorylation in vitro. *Environmental Toxicology*, 31(2): 245-255.
- Zhao, B., Yang, J., Liu, Z., Xu, Z., Qiu, Y., & Sheng, G. (2006). Joint anti-estrogenic effects of PCP and TCDD in primary cultures of juvenile goldfish hepatocytes using vitellogenin as a biomarker. *Chemosphere*, 65: 359-364.
- Zheng, W., Yu, H., Wang, X., & Qu, W. (2012). Systematic review of pentachlorophenol occurrence in the environment and in humans in China: not a negligible health risk due to the re-emergence of schistosomiasis. *Environ Int*, 42: 105-116.
- Zhou, L.Y., Wang, D.S., Shibata, Y., Bindhu, P.P., Suzuki, A., & Nagahama, Y. (2007). Characterization, expression and transcriptional regulation of P450c17-I and -II in the medaka, (*Oryzias latipes*). *Biochem Biophys Res Commun*, 362(3): 619-625.